

Appendix to: The level 1 weight 2 case of Serre's conjecture - a strategy for a proof

Luis V. Dieulefait
Universitat de Barcelona *

February 1, 2008

1 A fact, a question, and its answer

(we follow the notation of the preprint)

As explained in the preprint, to complete the proof our task is to show that, among the minimal (modular) Barsotti-Tate p -adic deformations of $\hat{\rho}|_{G_F}$ there is at least one that can be extended to $G_{\mathbb{Q}}$.

The field F is known to be unramified at p and we can choose it (as explained in the preprint, using solvable base-change and Sylow theorems) so that its degree over $\mathbb{Q}$ is prime to p .

With these conditions, as we already explained, there is a one-to-one correspondence between minimally ramified Barsotti-Tate deformations of $\hat{\rho}$ and those minimally ramified Barsotti-Tate deformations of $\hat{\rho}|_{G_F}$ that can be extended to the full $G_{\mathbb{Q}}$. (CC)

Let us call R the universal deformation ring of minimally ramified (Barsotti-Tate at p) deformations of $\hat{\rho}$, and let R' denote a similar minimal universal deformation ring, but of $\hat{\rho}|_{G_F}$.

We know from the results of Taylor that R' is a complete intersection ring.

*e-mail: ldieulefait@ub.edu

We will also need the following fact (proved by Bockle and Ramakrishna):

FACT (FF): Let W be the Witt ring of $\mathbb{F}_q$, then R is an W -algebra of the type:

$$W[[X_1, \dots, X_r]]/(f_1, \dots, f_s)$$

with $r \geq s$.

Now, let us answer the question: Why in the complete intersection ring R' there is at least one p -adic deformation that survives when we descend to $G_{\mathbb{Q}}$?

Well, if none of the minimal deformations of $\hat{\rho}|_{G_F}$ descends to $G_{\mathbb{Q}}$, then using the correspondence (CC) we see that R is too small to match with (FF).

This concludes the proof, so existence of minimal deformations, and Serre's conjecture in the level 1 weight 2 case follow.